



Effects of fermentation conditions and homogenization pressure on the rheological properties of Kefir

Meriem Bensmira, Consolate Nsabimana, Bo Jiang*

State Key Laboratory of Food Science and Technology, Jiangnan University, 1800 Lihu Avenue, 214122 Wuxi, Jiangsu, China

ARTICLE INFO

Article history:

Received 27 November 2009

Received in revised form

31 March 2010

Accepted 8 April 2010

Keywords:

Kefir

Fermentation temperature and time

Homogenization pressure

Rheology

EPS production

ABSTRACT

In this study, the effects of fermentation conditions (temperature and time) as well as homogenization pressure on the rheology and EPS production in Kefir made from bovine whole milk were investigated. Results showed that the rheological characteristics and EPS production are affected significantly ($p < 0.05$) by the fermentation temperature but not by the incubation time. Furthermore, the homogenization pressure was found to influence significantly ($p < 0.05$) the rheology but not the production of EPS in Kefir.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Fermented milks or beverages made with co-cultures of lactic acid bacteria and yeasts are widely produced in many countries in the region between Eastern Europe and Mongolia (Kurmann, Rašić, & Kroger, 1992). Some typical examples are Acidophilin, Acidophilus yeast beverage or milk, as well as Kefir and Koumiss. The latter product is traditionally made from mare's and/or yak's milk, whilst Kefir is mainly produced from bovine milk (Wszolek, Tamime, Muir, & Barclay, 2001). Kefir can be produced by fermenting milk with commercial freeze-dried Kefir starter cultures or traditional Kefir grains as well as the product obtained after the removal of grains. Kefir grains contain a wide and varying microflora, including lactic acid bacteria, acetic acid bacteria, yeasts and moulds (Pintado, Da Silva, Fernandes, Malcata & Hogg, 1996).

Gel formation is the most important physical property of fermented milk products. According to Tamime and Robinson (1989), rheological characteristics of milk gel are governed by milk composition, dry matter content, heating, homogenization, incubation temperature, cooling, storage time.

Studies on the rheological properties of Kefir are very scarce (Paraskevopoulou et al., 2003; Wszolek et al., 2001) and researches on the effects of processing conditions on Kefir quality are quite inexistent. The objective of this work was to make a contribution to

the knowledge of Kefir quality in terms of rheology and EPS production, with a particular focus on the effects of fermentation time and temperature as well as homogenization pressure.

2. Materials and methods

2.1. Milk types and Kefir starter culture

A commercial freeze-dried Kefir starter culture (coded BE010) was purchased from "Wilderness Family Naturals" (Silver Bay, Minnesota, USA). According to the supplier, this starter culture contains a mixture of *Lactococcus lactis* subsp. *lactis*, *L. lactis* subsp. *cremoris*, *L. lactis* subsp. *diacetylactis*, *Leuconostoc mesenteroides* subsp. *cremoris*, *Lactobacillus kefir*, *Kluyveromyces marxianus* var. *marxianus* and *Saccharomyces unisporus* (www.wildernessfamilynaturals.com). Skimmed milk powder (Guangming brand) and cow's raw milk were obtained from local supermarket and farm, respectively (Wuxi, China).

2.2. Preparation of Kefir working-culture

Sterile reconstituted skimmed milk was prepared by reconstituting approximately 12 g of skimmed milk powder in 100 ml of distilled water (Garrote, Abraham, & De Antoni, 1997), storing overnight at 4 °C to allow full rehydration, sterilizing at 121 °C for 2 min (Ozer, Robinson, Grandison, & Bell, 1998) and cooling down to about 25 °C. Reactivation was carried out by inoculating the sterile reconstituted skimmed milk with the commercial freeze-dried

* Corresponding author. Tel.: +86 510 85329055; fax: +86 510 85919625.

E-mail address: bjiang@jiangnan.edu.cn (B. Jiang).

culture at 0.2 mg/100 ml, incubating at 22 °C for 24 h (Beshkova, Simova, Frengova, Simov & Dimitrov, 2003) and ripening at 4 °C for 24 h. Kefir working-culture was prepared by inoculating the sterile reconstituted skimmed milk with the reactivated culture at 3 ml/100 ml, fermenting at 22 °C for 24 h (Beshkova et al., 2003) and maturing at 4 °C for at least one day before further use.

2.3. Kefir manufacture

Kefir was prepared as described by Tamime, Muir, and Wszolek (1999) with some modifications. Bovine farm milk was divided into three batches which were pre-heated to 60 °C, homogenized at 15 MPa (Tamime & Robinson, 1999) and pasteurized at 92 °C for 15 min (Beshkova, Simova, Simov, Frengova, & Spasov, 2002) using a water-bath. The pasteurized milk was first cooled down to about 25 °C and then inoculated with the working-culture at 3 ml/100 ml. The effect of fermentation time was investigated by incubating milk samples in a thermostatically controlled incubator at 22 °C (Beshkova et al., 2003) for various periods of time (18, 24 and 30 h). The impact of fermentation temperature was evaluated by incubating milk samples for 18 h at different temperatures (20, 24 and 28 °C). The influence of homogenization pressure was studied by dividing bovine farm milk into four batches and pre-heating milk samples to 60 °C. Only one batch was pasteurized directly without homogenization, while the other three batches were homogenized prior to pasteurization. Homogenization was carried out using APV-1000 homogenizer (APV Co., U.K.) in a single-stage (15, 17, 19 and 21 MPa). Milk samples were pasteurized, cooled down, and inoculated as mentioned above. Fermentation took place in a thermostatically controlled incubator at 24 °C for 18 h.

2.4. Rheological analysis

Dynamic rheological measurements were carried out using an AR 1000 rheometer (TA Instruments Ltd., Surrey, England) with a 1 mm-gap parallel-plate sensor. A portion of sample was carefully removed and placed on the bottom plate of the rheometer. The top plate was slowly lowered until the gap was 1 mm and excess sample was removed from the edges of the plate. Following the method of Rimada and Abraham (2006), two types of measurements were done: a strain sweep at constant frequency (1 Hz) to determine the linear viscoelasticity range and a frequency sweep from 0.0368 to 10.0 Hz at the same deformation (1%), within the linear range. The elastic modulus (G') and the viscous modulus (G'') were recorded as a function of frequency. The complex dynamic viscosity (η^*) was measured according to Haque, Richardson, and Morris (2001) by varying the frequency at a fixed strain of 1%. Kefir samples were prepared in triplicate for each batch and were analyzed individually.

2.5. Determination of EPS production

The EPS production was determined following the method of Purwandari, Shah, and Vasiljevic (2007). Results were expressed as the amount of crude EPS per gram of Kefir.

2.6. Statistical analysis

Results were treated using the analysis of variance (ANOVA) procedure of the SAS System for Windows V8 (SAS, 1999). Kefir samples were prepared in triplicate for each batch and were analyzed individually. Data were expressed as mean values \pm Standard Deviation (SD). The Duncan's Multiple-Range Test (DMRT) was used to compare means with 5% as significance level.

3. Results and discussion

3.1. Effect of fermentation time on Kefir rheology and EPS production

Fig. 1 shows the changes of storage modulus (G') and loss modulus (G'') as well as complex viscosity (η^*) values with fermentation time (1 Hz frequency and 1% strain). It is interesting to note that G' , G'' and η^* values increased gradually with the fermentation time. In fact, Kefir samples incubated at 22 °C for 18 and 30 h exhibited the lowest (269.40 Pa, 78.01 Pa and 38.36 Pa.s) and highest (548.76 Pa, 153.13 Pa and 77.92 Pa.s) values of G' , G'' and η^* , respectively. Although Kefir samples were significantly different ($p < 0.05$) in terms of G' , G'' and η^* , however no significant difference ($p > 0.05$) could be found among Kefir samples incubated at different time periods when other rheological characteristics such as $\tan \delta$ and G^* were taken into consideration (data not shown). In addition, the production of EPS seemed to increase in function of incubation time, with the lowest (421.66 $\mu\text{g/g}$) and highest (586.66 $\mu\text{g/g}$) amounts of EPS being observed for Kefir samples fermented at 22 °C for 18 and 24 h, respectively (Table 1). Nevertheless, Kefir samples were not significantly different from each other ($p > 0.05$).

The increase of G' , G'' and η^* during fermentation time can be attributed to the increase of metabolic activity of lactic acid bacteria, which promote the association of casein micelles by lowering pH. Certain strains of lactic acid bacteria make a further contribution to the physical structure of fermented milks (Rohm & Kovac, 1994; Vlahopoulou & Bell, 1993) by producing extracellular polysaccharides (EPS). According to Renard, van de Velde & Visschers (2006), the polysaccharides synthesized by lactic acid bacteria during fermentation increase the viscosity of the serum phase, bind hydration water and reduce consequently the water flow in the matrix space. In this study, the decrease in EPS production recorded after 30 h may be due to the presence of glycohydrolases capable of hydrolyzing EPS and liberating monomers. This observation coincides with that of Pham, Dupont, Roy, Lapointe, and Cerning (2000) as well as Lin & Chang Chien, (2007).

3.2. Effect of fermentation temperature on Kefir rheology and EPS production

Fig. 2 illustrates the evolution of storage modulus (G'), loss modulus (G'') and complex viscosity (η^*) in function of incubation temperature at 1 Hz frequency and 1% strain. The values of G' and η^* increased gradually with fermentation temperature while those of G'' showed a slight increment. Indeed, the lowest (199.46 Pa, 58.54 Pa and 28.43 Pa.s) and highest (790.00 Pa, 223.43 Pa and




Fig. 1. Changes of G' (\square), G'' (\blacksquare) and η^* (\blacksquare) at 1 Hz and 1% strain as a function of fermentation time in Kefir prepared from milk homogenized at 15 MPa and incubated at 22 °C. Each experiment was repeated three times and values are means \pm SD.




Fig. 4. Mechanical spectra of Kefir prepared using different conditions: (a) samples resulting from milk homogenization at 15 MPa and fermentation at 22 °C for 18 h; (b) samples obtained by homogenizing milk at 15 MPa and fermenting at 24 °C for 18 h; (c) samples corresponding to milk homogenized at 19 MPa and fermented at 24 °C for 18 h. G' (◆); G'' (▲) and η^* (○). Each experiment was repeated three times and values are means \pm SD.

homogenized at 15 MPa and fermented at 22 °C for 18 h, the storage (G') and loss (G'') moduli increased from 142.60 to 389.80 Pa and from 41.73 to 118.76 Pa, respectively while the complex viscosity (η^*) decreased from 642.60 to 6.52 Pa s with the increase of frequency (Fig. 4a). Similarly, Kefir sample made from milk homogenized at 15 MPa and fermented at 24 °C for 18 h exhibited G' , G'' and η^* values varying from 232.33 to 625.03 Pa, from 66.83 to 191.56 Pa and from 1045.33 to 10.46 Pa s, respectively within the frequency range used (Fig. 4b). Moreover, the values of G' , G'' and η^* in Kefir sample obtained after homogenizing milk at 19 MPa and fermenting it at 24 °C for 18 h ranged from 358.73 to 959.53 Pa, from 101.75 to 287.16 Pa and from 1612.33 to 16.03 Pa s, respectively in function of frequency (Fig. 4c). A decrease of the complex viscosity (η^*) with the increase of frequency has been reported also on naturally fermented milk by Narvhus, Østeraas, Mutukumira, and Abrahamsen (1998).

4. Conclusion

This study demonstrated that fermentation temperature and homogenization pressure are important processing parameters that can significantly ($P < 0.05$) affect the rheological properties of Kefir. However, the prolongation of fermentation time does not have a significant ($P > 0.05$) effect on the production of EPS in

Kefir. Consequently, the combination of a reasonable homogenization pressure (19 MPa), a moderate incubation temperature (24 °C) and a short fermentation time (18 h) have been found to be the optimal conditions for Kefir processing. However, a further investigation is needed to identify other factors susceptible to influence the quality characteristics of Kefir.

References

- Abbasi, H., Ehsani, M. R., Mousavi, S. M. A. E., D-Jomeh, Z. E., & Vaziri, M. (2007). Influence of exopolysaccharide producing starter cultures and incubation temperatures on the physical and rheological properties of low fat set type yogurt. Book of Abstracts (pp. 16–20). Copenhagen, Denmark: European Congress of Chemical Engineering (ECCE-6).
- Beshkova, D. M., Simova, E. D., Frengova, G. I., Simov, Z. I., & Dimitrov, Zh. P. (2003). Production of volatile aroma compounds by kefir starter cultures. *International Dairy Journal*, 13(7), 529–535.
- Beshkova, D. M., Simova, E. D., Simov, Z. I., Frengova, G. I., & Spasov, Z. N. (2002). Pure cultures for making kefir. *Food Microbiology*, 19(5), 537–544.
- Garrote, G. L., Abraham, A. G., & De Antoni, G. L. (1997). Preservation of kefir grains, a comparative study. *Lebensmittel-Wissenschaft Und-Technologie*, 30(1), 77–84.
- Haque, A., Richardson, R. K., & Morris, E. R. (2001). Effect of fermentation temperature on the rheology of set and stirred yogurt. *Food Hydrocolloids*, 15(4–6), 593–602.
- Küçükçetin, A., Weidendorfer, K., & Hinrichs, J. (2009). Graininess and roughness of stirred yoghurt as influenced by processing. *International Dairy Journal*, 19(1), 50–55.
- Kurmann, J. A., Rasić, J. I., & Kroger, M. (1992). *Encyclopedia of fermented fresh milk products*. New York: Van Nostrand Reinhold.
- Lankes, H., Ozer, H. B., & Robinson, R. K. (1998). The effect of elevated milk solids and incubation temperature on the physical properties of natural yoghurt. *Milchwissenschaft*, 53(9), 510–513.
- Lin, T. Y., & Chang Chien, M.-F. (2007). Exopolysaccharides production as affected by lactic acid bacteria and fermentation time. *Food Chemistry*, 100(4), 1419–1423.
- Narvhus, J. A., Østeraas, K., Mutukumira, T., & Abrahamsen, R. K. (1998). Production of fermented milk using a malty compound-producing strain of *Lactococcus lactis* subsp. *Lactis* biovar. *diacetylactis*, isolated from Zimbabwean naturally fermented milk. *International Journal of Food Microbiology*, 41(1), 73–80.
- Ozer, B. H., Robinson, R. K., Grandison, A. S., & Bell, A. E. (1998). Gelation properties of milk concentrated by different techniques. *International Dairy Journal*, 8(9), 793–799.
- Paraskevopoulou, A., Athanasiadis, I., Blekas, G., Koutinas, A. A., Kanellaki, M., & Kiosseoglou, V. (2003). Influence of polysaccharide addition on stability of a cheese whey kefir-milk mixture. *Food Hydrocolloids*, 17(5), 615–620.
- Peters, I. I. (1956). Cheddar cheese made from pasteurized milk homogenized at various pressures. *Journal of Dairy Science*, 39(8), 1083–1088.
- Pham, P. L., Dupont, I., Roy, G., Lapointe, G., & Cerning, J. (2000). Production of exopolysaccharide by *Lactobacillus rhamnosus* R and analysis of its enzymatic degradation during prolonged fermentation. *Applied Environmental Microbiology*, 66(6), 2302–2310.
- Pintado, M. E., Da Silva, J. A. L., Fernandes, P. B., Malcata, F. X., & Hogg, T. A. (1996). Microbiological and rheological studies on Portuguese kefir grains. *International Journal of Food Science & Technology*, 31(1), 15–26.
- Puhan, Z. (1988). Treatment of milk prior to fermentation. *International Dairy Federation Bulletin*, 227, 66–74.
- Purwandari, U., Shah, N. P., & Vasiljevic, T. (2007). Effects of exopolysaccharide-producing strains of *Streptococcus thermophilus* on technological and rheological properties of set-type yoghurt. *International Dairy Journal*, 17(11), 1344–1352.
- Renard, D., van de Velde, F., & Visschers, R. W. (2006). The gap between food gel structure, texture and perception. *Food Hydrocolloids*, 20(4), 423–431.
- Rimada, P. S., & Abraham, A. G. (2006). Kefiran improves rheological properties of glucono- δ -lactone induced skim milk gels. *International Dairy Journal*, 16(1), 33–39.
- Rohm, H., & Kovac, A. (1994). Effects of starter cultures on linear viscoelastic and physical properties of yogurt gels. *Journal of Texture Studies*, 25(3), 311–329.
- Ross-Murphy, S. B. (1984). Rheological methods. In H. W. -S (Ed.), *Chan, Biophysical methods in food research* (pp. 138–199). Oxford: Blackwell Scientific Publications.
- SAS. (1999). *SAS system for windows, version 8.01*. Cary, NC, USA: SAS Institute Inc.
- Sharma, R., Singh, H., & Taylor, M. W. (1996). Composition and structure of fat globule surface layers in recombined milk. *Journal of Food Science*, 61(1), 28–32.
- Tamime, A. Y., Muir, D. D., & Wszolek, M. (1999). Kefir, koumiss and kishk. *Dairy Industry International*, 64(5), 32–33.
- Tamime, A. Y., & Robinson, R. K. (1989). *Yogurt science and technology*. UK: Pergamon Press.
- Tamime, A. Y., & Robinson, R. K. (1999). *Yoghurt science and technology* (2nd ed.). New York: CRC Press.
- Vlahopoulou, I., & Bell, A. E. (1993). Effect of various starter cultures on the viscoelastic properties of bovine and caprine yogurt gels. *International Journal of Dairy Technology*, 46(2), 61–63.

- Vlahopoulou, I., Bell, A. E., & Wilbey, R. A. (2001). Effects of starter culture and its exopolysaccharides on the gelation of glucono- δ -lactone-acidified bovine and caprine milk. *International Journal of Dairy Technology*, 54(4), 135–140.
- van Vliet, T. (1988). Rheological properties of filled gels. Influence of filler matrix interaction. *Colloid and Polymer Science*, 266(6), 518–524.
- van Vliet, T., & Dentener-Kikkert, A. (1982). Influence of the composition of the milk fat globule membrane in the rheological properties of acid milk gels. *Netherlands Milk and Dairy Journal*, 36, 261–265.
- Wszolek, M., Tamime, A. Y., Muir, D. D., & Barclay, M. N. I. (2001). Properties of Kefir made in Scotland and Poland using bovine, caprine and ovine milk with different starter cultures. *Lebensmittel-Wissenschaft Und-Technologie*, 34(4), 251–261.